

1. Mohs Surgery is strongly indicated for which of the following?
 - a. Malignant Melanoma
 - b. Low grade Basal Cell Cancer
 - c. Recurrent cutaneous Squamous Cell Cancer
 - d. Mucosal Squamous Cell Cancer
2. What is the cure rate of Mohs surgery for primary basal cell cancer?
 - a. 69%
 - b. 79%
 - c. 89%
 - d. 99%
3. Which of the following is most amenable to secondary healing?
 - a. Helical rim defects
 - b. Nasal tip defects
 - c. Cheek defect
 - d. Forehead defect
4. Which of the following is a transposition flap?
 - a. Paramedian forehead flap
 - b. Melolabial flap
 - c. Rhomboid flap
 - d. Radial Forearm flap
5. Which of the following is true of eyelid defects?
 - a. Lower lid defects often heal well secondarily
 - b. Upper lid defects heal well secondarily
 - c. Lateral canthotomy is rarely required for large defect closure
 - d. None of the above
6. Which of the following is true of auricular defects?
 - a. Helical rim defects > 1.5cm can be closed primarily
 - b. If perichondrium is intact on one side of the cartilage, chance for favorable secondary healing is good
 - c. The lobule heals well secondarily
 - d. The conchal bowl heals poorly secondarily
7. What is the best flap to close a lower central lip defect 60% the length of the lower lip?
 - a. Wedge excision
 - b. M-plasty
 - c. Karapandzic Flap
 - d. Estlander Flap
8. What percentage of tumor margin is visualized with convention frozen section?
 - a. 100%
 - b. 80%
 - c. 25%
 - d. <1%
9. True or False: Mohs surgery is useful in Dermatofibrosarcoma
10. True or False: Nasal reconstruction should not excise more skin than the initial defect.