

■ **Diagnostic criteria for 299.00 Autistic Disorder**

- A. A total of six (or more) items from (1), (2), and (3), with at least two from (1), and one each from (2) and (3):
- (1) qualitative impairment in social interaction, as manifested by at least two of the following:
 - (a) marked impairment in the use of multiple nonverbal behaviors such as eye-to-eye gaze, facial expression, body postures, and gestures to regulate social interaction
 - (b) failure to develop peer relationships appropriate to developmental level
 - (c) a lack of spontaneous seeking to share enjoyment, interests, or achievements with other people (e.g., by a lack of showing, bringing, or pointing out objects of interest)
 - (d) lack of social or emotional reciprocity
 - (2) qualitative impairments in communication as manifested by at least one of the following:
 - (a) delay in, or total lack of, the development of spoken language (not accompanied by an attempt to compensate through alternative modes of communication such as gesture or mime)
 - (b) in individuals with adequate speech, marked impairment in the ability to initiate or sustain a conversation with others
 - (c) stereotyped and repetitive use of language or idiosyncratic language
 - (d) lack of varied, spontaneous make-believe play or social imitative play appropriate to developmental level

(continued)

□ **Diagnostic criteria for 299.00 Autistic Disorder** *(continued)*

- (3) restricted repetitive and stereotyped patterns of behavior, interests, and activities, as manifested by at least one of the following:
 - (a) encompassing preoccupation with one or more stereotyped and restricted patterns of interest that is abnormal either in intensity or focus
 - (b) apparently inflexible adherence to specific, nonfunctional routines or rituals
 - (c) stereotyped and repetitive motor mannerisms (e.g., hand or finger flapping or twisting, or complex whole-body movements)
 - (d) persistent preoccupation with parts of objects
- B. Delays or abnormal functioning in at least one of the following areas, with onset prior to age 3 years: (1) social interaction, (2) language as used in social communication, or (3) symbolic or imaginative play.
- C. The disturbance is not better accounted for by Rett's Disorder or Childhood Disintegrative Disorder.