

Section: UTMB On-line Documentation	01.34 - Policy
Subject: Healthcare Epidemiology Policies and Procedures	05.24.18 - Revised
Topic: 01.34 - Varicella-Zoster Virus Infection Control Program	1994 - Author

01.34 - Varicella-Zoster Virus Infection Control Program

Purpose	To describe a Varicella-Zoster Virus (VZV) Infection Control Program.
Audience	All employees of UTMB hospitals, clinics, Victory Lakes outpatient specialty care and surgical center, contract workers, volunteers and students with patient contact.
Policy	<p>It is the intent of this program to prevent the nosocomial transmission of VZV between patients and between patients and healthcare workers.</p> <p>All healthcare workers (HCWs) who have patient contact or work in patient care areas must have evidence of immunity (definite history of chickenpox, positive serologic test for VZV or two doses of VZV vaccine) to VZV. It shall be the responsibility of the employee health service or respective management group to assure that all healthcare workers who have patient contact or work in patient care areas are immune to VZV or have received two doses of the VZV vaccine at least 4 weeks apart.</p>
Employee Screening and Vaccination	<p>At the time of the new-hire employee health evaluation, proof of positive immunity to VZV will be established by a written documentation of VZV infection (chickenpox or herpes zoster) or VZV vaccination from a health care provider. Current employees may be screened for immunity to VZV as well.</p> <p>The proof of positive immunity to VZV shall be recorded in the employee's health record by Employee Health.</p> <p>A negative or equivocal history of VZV infection or vaccination shall be recorded in the employee's health record, and these employees shall be tested for VZV immunity by antibody test. If a serologic test for VZV antibody is negative and there are no contraindications to vaccination, the employee will be immunized with two doses of the VZV vaccine (see section on immunization).</p>

Section: UTMB On-line Documentation	01.34 - Policy
Subject: Healthcare Epidemiology Policies and Procedures	05.24.18 - Revised
Topic: 01.34 - Varicella-Zoster Virus Infection Control Program	1994 - Author

Immunization
with VZV
Vaccine

Individuals who are seronegative for VZV will be given the VZV vaccine, if they have none of the following contraindications for vaccination:

- History of an anaphylactic reaction to neomycin
- Presence of severe illness
- Presence of untreated active tuberculosis
- History of altered immunity
- Malignant neoplasms
- Primary (inherited) immunodeficiency including individuals with first degree relatives who have a primary immunodeficiency syndrome
- Acquired immunodeficiencies including HIV infection, cellular immunodeficiencies, hypogammaglobulinemia, dysgammaglobulinemias, and steroid therapy with $\geq 2\text{mg/kg}$ body weight per day
- Pregnancy
- Receipt of blood or plasma transfusions, immune globulin, or VaricellaZoster Immune Globulin (VariZIG) in the last 5 months

Vaccine shall be administered subcutaneously in two doses of 0.5 ml 4 to 8 weeks apart.

The vaccine must be stored at an average temperature of -15°C or $+5^{\circ}\text{F}$ or colder.

Vaccine must be used within 30 minutes of reconstitution. Vaccine remaining after this time shall be discarded.

Post-vaccination rash in employees.

- Employees will be instructed that they may develop a rash 2-6 weeks after vaccination and that this rash may be either a localized rash at the site of vaccination or a diffuse varicella-like rash. Either rash may be atypical with macules or papules rather than vesicles. Employees will be instructed to report immediately to the Employee Health Service if a rash develops.
- Employees with either an injection site rash or a generalized rash will be furloughed until the rash resolves (usually in 2-3 days).
- The Employee Health Service will verify that the rash has resolved and that the employee is not potentially infectious to others before allowing the employee to resume patient care duties.

A contact investigation shall not be done as the result of exposure to a healthcare worker with a post-vaccination skin rash.

Section: UTMB On-line Documentation	01.34 - Policy
Subject: Healthcare Epidemiology Policies and Procedures	05.24.18 - Revised
Topic: 01.34 - Varicella-Zoster Virus Infection Control Program	1994 - Author

Isolation for VZV Infection

Patients with suspected or documented primary VZV infection (chickenpox) and immunocompromised patients with localized or disseminated herpes zoster shall be placed on Contact and Airborne Precautions.

Immunocompetent patients with localized herpes zoster shall be placed on Contact Precautions.

Isolation shall continue until all skin lesions are crusted. Susceptible patients exposed to VZV shall be discharged from the hospital as soon as possible after notification of their attending physician.

Susceptible patients exposed to VZV infection who must remain hospitalized shall be placed on Contact and Airborne Precautions from 10 days after the first contact to 21 days after the last contact.

Exposure of Patients

Definition of exposure for patients (excluding perinatal exposure):

- Sharing contiguous air space with an infected patient or healthcare worker i.e., in the same room.
- Exposure must take place from 2 days before onset of rash to the time that all skin lesions are crusted in the index patient.

Perinatal exposure is defined as onset of primary VZV infection (chickenpox) in the mother from 5 days before birth until 2 days after birth.

Investigation and prevention for patients exposed to the Varicella-Zoster Virus:

- A history of primary VZV infection (chickenpox) or reactivation of latent VZV infection (herpes zoster or shingles) shall be taken from each exposed patient or their parent or guardian.
- Patients with the following conditions that place them at increased risk for severe VZV disease shall be identified and recorded:
 - Patients with immunocompromising diseases and treatments
 - Patients with underlying skin diseases and burns
 - Pregnancy
- Serologic tests shall be performed for detection of VZV antibody in the aforementioned patients regardless of prior VZV immunity history.
- Serologic tests shall be performed for detection of VZV antibody in all other patients with negative or equivocal histories for VZV infection.
- Patients without immunity to VZV shall be placed on Contact and Airborne Precautions from 10 days after the first exposure until 21 days after the last exposure (28 days for patients who received VariZIG, see below).

Exposure of Healthcare Workers

Definition of exposure for healthcare workers:

- Sharing contiguous air space with an infected patient or healthcare worker in the same room.
- Exposure must take place from 2 days before onset of rash to the time that all skin lesions are crusted in the index patient.

Section: UTMB On-line Documentation	01.34 - Policy
Subject: Healthcare Epidemiology Policies and Procedures	05.24.18 - Revised
Topic: 01.34 - Varicella-Zoster Virus Infection Control Program	1994 - Author

Investigation and prevention for HCWs exposed to the Varicella-Zoster Virus:

- A history of prior VZV immunity shall be taken from each exposed healthcare worker. Employee's records shall be checked by Employee Health Clinic staff for evidence of a positive serologic test and previous vaccinations with VZV vaccine.
- Serologic tests shall be performed for detection of VZV antibody for healthcare workers with a negative or equivocal history for VZV immunity who have not been vaccinated
- Healthcare workers who have no positive immunity to VZV documented in Employee Health records, shall also be given prophylaxis (see Prophylaxis section).
- Healthcare workers who have underlying severe immunocompromising diseases, or are on immunocompromising therapy VZV may also be offered prophylaxis.
- Healthcare workers who have positive proof of prior VZV immunity as recorded in Employee Health records may continue to work with patients and do not require prophylaxis.

Acyclovir/
Valacyclovir
Prophylaxis

Acyclovir and valacyclovir are class B antiviral drugs and are the preferred agents for prophylaxis in exposed persons.

Prophylaxis should be started 9 days after the first exposure and be given for seven days.

Prophylaxis for healthcare workers and adult patients:

- For adults weighing greater than 40 kgs, 1 gram of valacyclovir given orally 3 times per day.
- For adults weighing less than 40 kgs, 500 mg of valacyclovir given orally 3 times per day.

Prophylaxis for children:

- Acyclovir 80 mg/kg/day divided into 4 doses po.
- The maximum dose of acyclovir is 800 mg, 4 times per day.

Patients who receive acyclovir/valacyclovir prophylaxis and remain in the hospital will be placed on Airborne and Contact Precautions from 10 days after the first exposure until 21 days after the last exposure.

Healthcare workers who receive acyclovir/valacyclovir prophylaxis will be furloughed from 10 days after the first exposure until 21 days after the last exposure.

70% of susceptible persons who receive acyclovir (valacyclovir) prophylaxis and develop no clinical signs of chickenpox, will have antibodies to Varicella Zoster three weeks after completion of prophylaxis.

Section: UTMB On-line Documentation	01.34 - Policy
Subject: Healthcare Epidemiology Policies and Procedures	
Topic: 01.34 - Varicella-Zoster Virus Infection Control Program	05.24.18 - Revised
	1994 - Author

- All patients and healthcare personnel who receive acyclovir (valacyclovir) prophylaxis and do not develop chickenpox will have a serologic test for varicella done 3 weeks after completion of the course of prophylaxis.
- Healthcare workers who are seronegative will be given two doses of varicella-zoster vaccine 4 to 8 weeks apart.

Administration of
VariZIG

VariZig is very expensive and may have a lower efficacy than acyclovir/valacyclovir, and is therefore to be used ONLY for those exposed persons who have renal failure or proven previous allergy to acyclovir/valacyclovir. Approval for VariZig should be obtained from the staff of Healthcare Epidemiology (24/7 pager 409-643-3133).

VariZIG shall be administered as soon as possible after exposure but no later than 96 hours after the first exposure to VZV infection

VariZIG is administered intramuscularly in a dose of 125 units/10 kg (22lbs) of body weight up to a maximum of 625 units (i.e., 5 vials). The minimum dose is 125 units. Children who weigh less than 10 kgs should receive 125 units.

VariZIG SHALL NOT BE ADMINISTERED INTRAVENOUSLY.

After receiving VariZIG, patients must be kept in isolation for 28 days after the last exposure and healthcare workers must remain on furlough for 28 days after their last exposure.

Section: UTMB On-line Documentation	01.34 - Policy
Subject: Healthcare Epidemiology Policies and Procedures	05.24.18 - Revised
Topic: 01.34 - Varicella-Zoster Virus Infection Control Program	1994 - Author

References

1. Guideline for Infection Control in Healthcare personnel 1998 AJIC June 1998. 289-327.
2. Centers for Disease Control and Prevention. Prevention of Varicella: updated recommendations of the Advisory Committee on immunization practices (ACIP). MMWR 1999; 48 (No. RR-6): 1-4.
3. ACIP provisional recommendations for prevention of Varicella. Available at [http://www.cdc.gov/nip/vaccine/varicella/varicella_acip_recs_prov_june_2006.p df](http://www.cdc.gov/nip/vaccine/varicella/varicella_acip_recs_prov_june_2006.pdf). Accessed 05/20/07.
4. The Medical Letter. *VariZIG* for prophylaxis after exposure to Varicella. 2006; 48:69-70.
5. MMWR. A new product (*VariZIG*TM) for postexposure prophylaxis for Varicella available under an investigational new drug application expanded access protocol. 2006; 55(08): 209-210.
6. Chaves SS, Gargiullo P, Zhang JX, et al. Loss of Vaccine-induced immunity to Varicella over time. N Engl J Med 2007; 356:1121-1129.
7. CDC. National Immunization Program. Varicella Vaccine – FAQs about healthcare workers. Available at <http://www.cdc.gov/nip/vaccine/varicella/faqsclinic-vac-hcw.htm#5-rash> Accessed 05/30/07.
8. Huang Y-C, Lin T-Y, Chiu C-H. Acyclovir prophylaxis of Varicella after household exposure. Pediatr Infect Dis J. 1995; 14:152-154.
9. Lin T-Y, Huang Y-C, Ning H-C, Hsueh. Oral acyclovir prophylaxis of Varicella after intimate contact. Pediatr Infect Dis J 1997; 16:1162-1165.
10. Kumagai T, Kamada M, Igarash C, et al. Varicella-Zoster virus-specific cellular immunity in subjects given acyclovir after household chickenpox exposure. J Infect Dis 1999; 180:834-837.
11. Weinstock DM, Boeckh M, Boulad F, et al. Postexposure prophylaxis against varicella-zoster virus infection among recipients of hematopoietic stem cell transplant: unresolved issues. Infect Control Hosp Epidemiol 2004; 25:603608.
12. American Academy of Pediatrics. Red Book. 2006:716-717.