

<b>Section:</b>	<b>UTMB On-line Documentation</b>	<b>01.46-Policy</b>
<b>Subject:</b>	<b>Infection Control &amp; Healthcare Epidemiology Policies and Procedures</b>	<b>10.4.24-Revised</b>
<b>Topic:</b>	<b>01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)</b>	<b>2007-Author</b>

## 01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)

Purpose: Catheter associated urinary tract infections (CAUTIs) are prevented by the appropriate use of catheters and should never be used solely for the convenience of patient-care personnel (i.e. monitoring urine output) or patient preference (i.e. does not want to use urinal or immobile). The policy is intended to provide infection control guidelines for the proper placement and management of indwelling urinary catheter devices.

Audience: All employees of UTMB hospitals, clinics, outpatient surgical center, licensed independent practitioners, contract workers, and students.

- I. Catheter Use
  - A. Urinary catheters should be inserted only when necessary and left in place only for as long as necessary. They should not be used solely for the convenience of patient-care personnel or patient preference.
    1. Alternatives to indwelling catheters must be considered first if suitable in a specific patient. These include the use of external male ('condom') catheter and vacuum-assisted female and male catheters, clean intermittent catheterization (CIC) and bladder massage.
  - B. To avoid urethral strictures, urethral erosion associated with chronic indwelling urethral catheterization, suprapubic catheterization should be considered in patients who need prolonged bladder catheterization for more than 4 weeks (e.g. those with neurogenic bladder, chronic urethral obstruction or ulceration in perineal area). Urinary tract infection associated with suprapubic catheter is not reportable to CDC/NHSN (only transurethral catheter infections are reportable). Consultation with the urology team should be obtained for suprapubic catheter placement plan in either inpatient or outpatient settings.
- II. Leadership for Appropriate Catheter Use
  - A. The clinical unit dyads will oversee and support the safe use of urinary catheters as outlined in this policy.
- III. Indications for Indwelling Transurethral Catheters
  - A. Indwelling transurethral catheters must be inserted only when there is an indication to do so. Indications include the following but only when an external catheter cannot be used:
    1. Acute urinary retention.
      - a. A urethral catheter should be considered for the purpose of acute urinary retention when at least one (1) of the following criteria are met:
        - i. Clean intermittent catheterization has been done every 4-6 hours for 24 hours and post residual volume has remained above 300 mL.
        - ii. If self-voiding has not resumed within 48 hours and intermittent catheterization cannot be continued due to staffing or patient care issues, the urology team should be consulted for diagnosis and advice regarding the duration of catheter use as well as alternate urinary drainage methods.
      - b. Transurethral urinary catheter inserted for management of acute urinary retention should not be used for more than 48 hours. Patient should be re-evaluated for ability to self-void or resuming intermittent catheterization.
    2. Known or suspected chronic urinary retention or obstruction (tumors, strictures,

<b>Section:</b>	<b>UTMB On-line Documentation</b>	<b>01.46-Policy</b>
<b>Subject:</b>	<b>Infection Control &amp; Healthcare Epidemiology Policies and Procedures</b>	<b>10.4.24-Revised</b>
<b>Topic:</b>	<b>01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)</b>	<b>2007-Author</b>

- clots)
3. End of life care, IF needed (patient should not be undergoing tests and treatments other than pain control)
  4. Critically ill-need accurate I/O measurement
  5. Selected surgical procedures-GU or colorectal surgery
  6. Urinary diversion e.g: to assist in healing open sacral or perineal wound in the incontinent patient, Open wounds/ pressure ulcers stage 3 or 4 or unstageable sacral pressure ulcer IF cannot be kept clean with wound dressing or alternative urine collection devices
  7. Intraoperative monitoring
  8. Prolonged immobilization
- B. Orders for insertion and discontinuation
1. Urethral (Foley) catheters may be inserted in patients only by an order from a physician.
  2. When a urethral catheter is ordered by a physician, the physician must check the indication for catheter insertion from a drop-down list of indications in EPIC EMR.
  3. The initial insertion order will include the approval to discontinue the catheter based on nursing assessment and discussion with the primary care team.
  4. The nurse will assess the indications for a urinary catheter during each shift and will document in EPIC. The nurse will discontinue the catheter if the indications are not met unless the patient's primary physician team determines that the catheter is needed. In that case, the treating physician will document in patient's progress note the indication(s) and rationale and duration for continued use of the indwelling catheter. **Please see Appendix 2: UTMB Indwelling Urinary Catheter Nurse Driven Discontinuation Algorithm.**
- IV. Indwelling Transurethral Catheters Present on Admission or Placed Emergently
- A. If an indwelling urethral catheter is present on admission, it should be removed **immediately if contraindications do not exist (after review with the physician)**, or a new catheter inserted if still warranted. Consider alternatives, including external male and female urinary catheters.
  - B. If an Indwelling urethral catheter is placed emergently, it must be removed as soon as possible (**within 48 hours**) since adherence to aseptic technique cannot be ensured, a base line urine culture obtained, and a new catheter inserted if still warranted.
- V. Catheter Insertion Technique
- A. Personnel who insert urinary catheters must be trained in proper insertion technique.
  - B. Hand hygiene must be performed with an antimicrobial soap and water or an alcohol handrub before insertion and immediately before and after any manipulation of the catheter site or drainage system.
  - C. Catheters shall be inserted using aseptic technique and sterile equipment.
  - D. Sterile gloves, drape, sponges, and appropriate antiseptic solution for periurethral cleansing, and a single-use packet of sterile lubricant jelly shall be used for insertion.
  - E. As small a catheter as possible, consistent with good drainage, should be used to minimize urethral trauma.
  - F. Only one attempt at insertion is allowed for each catheter; a new catheter must

<b>Section:</b>	<b>UTMB On-line Documentation</b>	<b>01.46-Policy</b>
<b>Subject:</b>	<b>Infection Control &amp; Healthcare Epidemiology Policies and Procedures</b>	<b>10.4.24-Revised</b>
<b>Topic:</b>	<b>01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)</b>	<b>2007-Author</b>

be used for each attempt until the catheter can be inserted without contamination.

- G. Indwelling catheter should be properly secured after insertion to prevent movement and urethral traction and subsequent erosion, make sure it is without traction.

#### VI. Documentation for Catheter Insertion

- A. The following information must be documented in the patient's medical record after catheter insertion
  - 1. Indication(s) for catheter insertion
  - 2. Date and time of catheter insertion
  - 3. Individual who inserted the catheter
  - 4. The size, type of urethral catheter (silicone vs latex; straight vs Coude), any difficulty, and how many mL filled the balloon.
- B. Include documentation in the nursing flow sheet, nursing notes or physician orders.
- C. Documentation should be accessible in the patient's medical record and recorded in a standard format for data collection and quality improvement purposes.

#### VII. Closed Sterile Drainage

- A. A sterile, continuously closed drainage system sealed to the catheter must be maintained.
- B. If breaks in aseptic technique, disconnection, or leakage occur, the catheter and collecting system sealed to the catheter should be replaced using aseptic technique.

#### VIII. Irrigation

- A. Bladder irrigation through urethral catheter should be avoided unless continuous or manual bladder irrigation is ordered by a physician. Irrigation is typically done because of hematuria or blood clots and may be indicated following interventions such as surgery, a traumatic urinary catheter insertion, or complex radiation cystitis. Bladder irrigation is not indicated when a catheter is blocked by sediment; instead, the catheter should be replaced. Routine bladder irrigation also should not be done for prevention of infection.
- B. Bladder should be irrigated using the same aseptic precautions used for insertion of urethral catheter. The catheter-tubing junction must be disinfected before disconnection.

#### IX. Specimen Collection

- A. If small volumes of fresh urine are needed for examination, The sampling port should be cleansed with alcohol. After the alcohol has dried, urine should be aspirated with a sterile needle and syringe.
- B. Larger volumes of urine for special analyses should be obtained aseptically from the drainage bag.
- C. Culture collection
  - 1. Prior to obtaining urine culture, please refer to Appendix 3 (**Urine Culture Collection Algorithm**) for proper indication and stepwise investigative approach through urinalysis followed by culture.
  - 2. If the indwelling urinary catheter has been in place for at least 7 days, it should be removed, and a new catheter be inserted aseptically prior to

<b>Section:</b>	<b>UTMB On-line Documentation</b>	<b>01.46-Policy</b>
<b>Subject:</b>	<b>Infection Control &amp; Healthcare Epidemiology Policies and Procedures</b>	<b>10.4.24-Revised</b>
<b>Topic:</b>	<b>01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)</b>	<b>2007-Author</b>

- collection of urine sample for culture.
  - 3. Urinary catheter tips should not be cultured and are not acceptable for diagnosis of a urinary tract infection.
  - 4. Urine cultures must be obtained using appropriate technique, such as clean catch collection or catheterization. Specimens taken from an indwelling catheter must be aspirated from a disinfected sample port.
  - 5. In infants, urine specimens should be collected by catheterization or suprapubic aspiration; positive urine cultures from bag specimens are not acceptable.
  - 6. Urine specimens collected for culture will be sent to the laboratory in a tube with a boric acid preservative (gray or yellow-top tube).
- X. Urinary Flow and Collection Bag
- A. Unobstructed flow should be maintained.
  - B. To achieve free flow of urine:
    - 1. the catheter and collection tubing should be kept from kinking.
    - 2. the collection bag should be emptied regularly using a separate collection container for each patient (the drainage spigot and nonsterile collection container should never come in contact);
    - 3. collection bags should always be kept below the level of the bladder but should never touch the floor. Catheter should be secured using fixation devices without traction.
    - 4. If the catheter becomes obstructed, it should be flushed if it is new ( $\leq 48$  hours) with sterile normal saline or water (60-120 cc). A physician order is required for this procedure. If the procedure fails, then consider inserting a new catheter using the same aseptic technique described above. Catheters that are older than 48 hours should be removed and a new catheter be inserted. The newly inserted catheter must be sealed to a new sterile closed drainage system. For procedures involving bladder irrigation, see section IX above.
- XI. Routine Bathing and Perineal Care
- A. The perineum should be cleaned daily and as needed with soap and water and dried followed by an application of 2% chlorhexidine gluconate to reduce colonization of the perineal skin by bacteria as needed (e.g. cleaning after bowel movements).
  - B. Routine hygiene (e.g., cleaning of the perimeatal surface during daily bathing) is appropriate.
  - C. Patients will be bathed daily and as needed (e.g. after bowel movements) with chlorhexidine gluconate from navel to knees while the indwelling urinary catheter is in place. **Refer to Appendix 4 for CHG bathing guidelines.**
- XII. Catheter Change Interval: Indwelling urethral catheters should be changed only as clinically indicated.
- XIII. Use of Bladder Scanners
- A. Refer to Appendix 1 for Bladder Scan Protocol.
  - B. Nursing staff must be trained in their use.
  - C. The equipment must be adequately cleaned and disinfected between patients according to the manufacturer's instructions for use.
  - D. Use a portable bladder scanner to assess urine volume in patients undergoing intermittent catheterization to reduce unnecessary catheter insertions.
  - E. Bladder ultrasound readings should be taken immediately after voiding to get a

<b>Section:</b>	<b>UTMB On-line Documentation</b>	<b>01.46-Policy</b>
<b>Subject:</b>	<b>Infection Control &amp; Healthcare Epidemiology Policies and Procedures</b>	<b>10.4.24-Revised</b>
<b>Topic:</b>	<b>01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)</b>	<b>2007-Author</b>

more accurate assessment of residual volume.

- F. Each bladder ultrasound should be confirmed with a second reading.
- G. Consecutive readings should be taken until a full view of the bladder is obtained on the scanner.


XIV. Outcome Measures: Definitions of the National Healthcare Safety Network (NHSN) will be utilized to identify catheter-associated urinary tract infections.  
<http://www.cdc.gov/nhsn/pdfs/pscmanual/7psccauticurrent.pdf>

XV. Insertion of Urethral catheters in the Emergency Department (ED)

- A. All urethral catheters inserted in the ED must be ordered by a physician.
  - 1. When the order is entered, it must state the indication for insertion of a Urethral catheter.
  - 2. The only indications for Urethral catheters are listed in section IIIA.
  - 3. Urethral catheters are not indicated for:
    - a. Fall prevention
    - b. Routine urine specimens
    - c. Staff request
    - d. Excoriated skin
    - e. Altered mental status
- B. Alternatives to indwelling Urethral catheters
  - 1. Unisex urinals may be used by both male and female patients to avoid use of a Urethral catheter.
  - 2. External male and female urinary catheters should be used when possible.
  - 3. Bladder scanners should be used to measure post-void residuals rather than straight catheter insertions.
- C. Catheter insertion technique
  - 1. Urethral catheters may not be inserted by nursing students, medical students or untrained residents unless they are supervised by trained Registered Nurses.
  - 2. For patients who are obese, two nurses should work together to expose the meatus for safe insertion of Urethral catheters.
  - 3. The patient's genital region must be thoroughly cleaned with soap and water followed by application of chlorhexidine gluconate.
  - 4. Catheters must be inserted using aseptic technique and sterile equipment.
    - a. Sterile gloves, drape, sponges, and appropriate aseptic solution for periurethral cleansing, and a single-use packet of sterile lubricant jelly should be used for insertion.
    - b. As small a catheter as possible, consistent with good drainage, should be used to minimize urethral trauma.
    - c. Only one attempt at insertion is allowed for each catheter; a new catheter must be used for each attempt until the catheter can be inserted without contamination.
    - d. Indwelling catheters should be properly secured after insertion to prevent movement and urethral traction.


<b>Section:</b>	UTMB On-line Documentation	<b>01.46-Policy</b>
<b>Subject:</b>	Infection Control & Healthcare Epidemiology Policies and Procedures	10.4.24-Revised
<b>Topic:</b>	01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)	2007-Author

**APPENDIX 1: BLADDER SCAN PROTOCOL AFTER URETHRAL CATHETER REMOVAL**



<b>Section:</b>	UTMB On-line Documentation	<b>01.46-Policy</b>
<b>Subject:</b>	Infection Control & Healthcare Epidemiology Policies and Procedures	10.4.24-Revised
<b>Topic:</b>	01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)	2007-Author


**APPENDIX 2: UTMB Indwelling Urinary Catheter—Nurse Driven Catheter Discontinuation Algorithm**



<b>Section:</b>	UTMB On-line Documentation	<b>01.46-Policy</b>
<b>Subject:</b>	Infection Control & Healthcare Epidemiology Policies and Procedures	10.4.24-Revised
<b>Topic:</b>	01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)	2007-Author

**APPENDIX 3:**

**Urine Culture Collection Algorithm**





<b>Section:</b>	<b>UTMB On-line Documentation</b>	<b>01.46-Policy</b>
<b>Subject:</b>	<b>Infection Control &amp; Healthcare Epidemiology Policies and Procedures</b>	
<b>Topic:</b>	<b>01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)</b>	<b>11.3.23-Revised 2007-Author</b>

**APPENDIX 4:**

## Navel to Knees with Chlorhexidine Gluconate (CHG): Preventing Catheter-Associated Urinary Tract Infections (CAUTI)

### What is the “Navel to Knees with CHG” Campaign?

Extending catheter care and peri-care from the navel to knees with CHG wipes to prevent bacteria from migrating towards the catheter.

**Supplies & Frequency:**

- Clean pair of Gloves and 6 CHG wipes per each treatment.
- Daily and as needed (e.g. cleaning after bowel movements).



# 5 STEPS IN “NAVEL TO KNEES WITH CHG” TREATMENT

**1** Perform hand hygiene and don clean gloves.



**2** Use wipe #1 to clean the indwelling urinary catheter tubing.

- Start with the tubing at the urethra and wipe down the tubing at the urethra and wipe down the tubing to include the bifurcated tubing at the statlock device.



**3** Use wipe #2 to perform perineal care on one side.

*Female anatomy:* wipe labia away from urinary meatus down towards the anus.  
*Male anatomy:* wipe away from the urethra along the shaft of one side of the penis, then down along the perineum on the same side towards the anus.

Repeat action on the other side with wipe #3.



**4** Use wipe #4 to clean the abdomen and abdominal folds - wiping up in a zigzag (right to left) pattern away from the groin area to the navel.



**5** Use wipe #5 to clean one upper thigh leg - wiping down and away from the groin.

Repeat with wipe #6 to clean the other upper thigh.



Discard Gloves & CHG wipes. Perform hand hygiene.

Document NTK CHG treatment in I/O flowsheet within the Urethral Catheter LDA assessment

Urethral Catheter	
Urethral Catheter Properties	
Daily review of necessity	
Urine Catheter Maintenance	PerinealCath
Urine color	
Urine description	
Output (mL)	



Schmuckli, Y., Olson-Sisk, K., Rivett, J., & Charney-Barish, J. (2018). Navel to Knees With Chlorhexidine Gluconate. *Dimensions of Critical Care Nursing*, 38(5), 236-242.

<b>Section:</b>	<b>UTMB On-line Documentation</b>	<b>01.46-Policy</b>
<b>Subject:</b>	<b>Infection Control &amp; Healthcare Epidemiology Policies and Procedures</b>	
<b>Topic:</b>	<b>01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)</b>	<b>11.3.23-Revised 2007-Author</b>

## **References**

1. Agency for Healthcare Research and Quality, Rockville, MD. <https://www.ahrq.gov/hai/universal-icu-decolonization/universal-icu-ape3.html> Internet Citation: Appendix E. Training and Educational Materials (continued). Content last reviewed September 2013.
2. Apisarnthanarak A, Thongphubeth K, Sirinvaravong S, et al. Effectiveness of multifaceted hospitalwide quality improvement programs featuring an intervention to remove unnecessary urinary catheters at a tertiary care center in Thailand. *Infect Control Hosp Epidemiol* 2007; 28:791-798.
3. Carter EJ, Pallin DJ, Mandel L, et al. Emergency department catheter-associated urinary tract infection prevention: multisite qualitative study of perceived risks and implemented strategies. *Infect Control Hosp Epidemiol* 2015;
4. Cassir N, Thomas G, Hraiech S, et al. Chlorhexidine daily bathing: impact on healthcare-associated infections caused by gram-negative bacteria. *Am J Infect Control*. 2015;43:640-643.
5. Centers for Disease Control and Prevention. Healthcare Infection Control Practices Advisory Committee. Guideline for prevention of catheter-associated urinary tract infections 2009;Division of Healthcare Quality Promotion. 1-67
6. Chenoweth C, Saint S, Preventing catheter-associated urinary tract infections in the intensive care unit. *Crit Care Clin* 2013;29:19-32.
7. Dromerick AW, Edwards DF. Relation of postvoid residual to urinary tract infection during stroke rehabilitation. *Arch Phys Med and Rehabil* 2003; 84:1369-1372.
8. Fakh MG, Heavens M, Grotenmeyer J, et al. Avoiding potential harm by improving appropriateness of urinary catheter use in 18 Emergency Departments. *Ann Emerg Med* 2014;63:671-768.
9. Huang W-C, Wann S-R, Lins-L, et al. Catheter-associated urinary tract infections in intensive care units can be reduced by prompting physicians to remove unnecessary catheters. *Infect Control Hosp Epidemiol* 2004; 25:974-978.
10. Kassakian SZ, Mermel LA, Jefferson JA, et al. Impact of chlorhexidine bathing on hospital-acquired infections among general medical patients. *Infect Control Hosp Epidemiol*. 2011;32:238-243.
11. Knoll BM, Wright D, Ellingson L, et al. Reduction in inappropriate urinary catheter use at a Veterans Affairs Hospital through a multifaceted quality improvement project. *Clin Infect Dis* 2011;52:1283-1290.
12. Liang SY, Theodoro DL, Schuur JD, Marschall J. Infection Prevention in the Emergency Department. *Ann Emerg Med* 2014;64:299-313.
13. Lo E, Nicolle L, Classen D, et al. Strategies to prevent catheter-associated urinary tract infections in acute care hospitals. *Infect Control Hosp Epidemiol* 2008; 29 (Suppl 1):S41-S50.
14. Marra AR, Camargo TZS, Goncalves P, et al. Preventing catheter-associated urinary tract infection in the zero tolerance era. *Am J Infect Control* 2011; 39:817-822.
15. Meddings J, Rogers MA, Krein SL, et al. Reducing unnecessary urinary catheter use

<b>Section:</b>	<b>UTMB On-line Documentation</b>	<b>01.46-Policy</b>
<b>Subject:</b>	<b>Infection Control &amp; Healthcare Epidemiology Policies and Procedures</b>	
<b>Topic:</b>	<b>01.46 - Prevention of Catheter-Associated Urinary Tract Infections (CAUTI)</b>	<b>11.3.23-Revised 2007-Author</b>

and other strategies to prevent catheter-associated urinary tract infection: an integrative review. *BMJ Qual Saf* 2014;23:277-289.

16. Phillips JK. Integrating bladder ultrasound into a urinary tract infection-reduction project. *Am J Nurs* 2000;suppl:3-12.
17. Reilly L, Sullivan P, Ninni S, et al. Reducing Foley catheter device days in an intensive care unit. Using the evidence to change practice. *AACN Adv Crit Care*. 2006; 17:272-283.
18. Saint S, Kowalski CP, Forman J, et al. A multicenter qualitative study on preventing hospital-acquired urinary tract infection in U.S. hospitals. *Infect Control Hosp Epidemiol* 2008; 29:333-341.
19. Schmudde, Y., Olson-Sitki, K., Bond, J., & Chamberlain, J. (2019). Navel to Knees With Chlorhexidine Gluconate. *Dimensions of Critical Care Nursing*, 38(5), 236–240. <https://doi.org/10.1097/dcc.0000000000000371>
20. Schuur JD, Chambers JG, Hou PC. Urinary catheter use and appropriateness in U.S. Emergency Departments, 1995-2010. *Acad Emerg Med* 2014;21:292-300.
21. Scott RA, Oman KS, Makic MBF, et al. Reducing indwelling urinary catheter use in the Emergency Department: A successful quality-improvement initiative *J Emerg Nurs* 2014;40:237-244.
22. Stevens E. Bladder ultrasound: avoiding unnecessary catheterizations. *MEDSURG Nursing* 2005; 14:249-253.